

ASK YOUR KIDS CARE FOR YOUR KIDS EMBRACE YOUR KIDS



See Reverse for Questions that Can Save a Life

Sponsored by the SPC Regional Prevention Coalition

Always ask questions 1 and 2.	Past	Month
 Have you wished you were dead or wished you could go to sleep and not wake up? 		
2) Have you actually had any thoughts about killing yourself?		
If YES to 2, ask questions 3, 4, 5 and 6. If NO to 2, skip to question 6.		
3) Have you been thinking about how you might do this?		
4) Have you had these thoughts and had some intention of acting on them?	High Risk	
5) Have you started to work out or worked out the details of how to kill yourself? Did you intend to carry out this plan?	High Risk	
Always Ask Question 6	Life- time	Past 3 Months
6) Have you done anything, started to do anything, or prepared to do anything to end your life? <i>Examples</i> : Took pills, tried to shoot yourself, cut yourself, tried to hang yourself, took out pills but didn't swallow any, held a gun but changed your mind or it was		High Risk



If YES to 2 or 3, seek behavioral healthcare for further evaluation. If the answer to 4, 5 or 6 is YES, get <u>immediate help</u>: Call or text 988, call 911 or go to the emergency room. <u>STAY WITH THEM</u> until they can be evaluated.



Columbia Protocol app